The long journey to the Higgs boson and beyond at the LHC

Graduiertenkolleg 1504/2
Humboldt-Universität zu Berlin
5th November 2013



Drawing by Sergio Cittolin

Peter Jenni, Freiburg and CERN

The Large Hadron Collider project is a global scientific adventure, combining the accelerator, a worldwide computing grid and the experiments, initiated almost 30 years ago



It is a great privilege and pleasure to present now physics results from the first three years of operation


History of the Universe
pp physics at the LHC corresponds to conditions around here
BIG BANG
Inflation
possible dark matter relicts
cosmic microwave radiation visible
t 10-44 10-37 s
T 1032 1028
E 1019 1015
10-10 s
1015
102
10-5 s
1012
10-1
102 s
109
10-4
3x105 y
3000
3x10-10
109 y
15
10-12
Today
12x109 y
2.7
2.3x10-13
(sec,yrs)
(Kelvin)
(GeV)
Key:
q quark
g gluon
e electron
m muon t tau
n neutrino
W, Z bosons
meson
baryon
ion
atom
photon
star
galaxy
black hole
HI physics at the LHC corresponds to conditions around here
Particle Data Group, LBNL, © 2000. Supported by DOE and NSF



Study physics laws of first moments after Big Bang
increasing Symbiosis between Particle Physics, Astrophysics and Cosmology

The Standard Model of Particle Physics





(i) Constituents of matter: quarks and leptons

(ii) Four fundamental forces
(described by quantum field theories, except gravitation)

(iii) The Higgs field (problem of mass)

Standard Model of Elementary Particles



On Tuesday 8th October 2013:







"for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at CERN's Large Hadron Collider"

Unification of Forces



The SM is not a complete theory

Some of the outstanding questions in fundamental physics addressed, at least in part, with the LHC are:

What is the origin of the elementary particle masses ? (~✔)

What is the nature of the Universe dark matter ?

Why is only matter observed in the Universe as primary constituents and not anti-matter ?

What are the features of the primordial plasma present ~10 μs after the Big Bang ?

What happened in the first moments of the Universe ~10^{-11} s after the Big Bang ?

Are there other forces in addition to the known four ?
Are there additional (microscopic) space dimensions ?
….

The SM is not a complete theory

Some of the outstanding questions in fundamental physics addressed, at least in part, with the LHC are:

What is the origin of the elementary particle masses ? (~✔)

What is the nature of the Universe dark matter ?

Why is only matter observed in the Universe as primary constituents and not anti-matter ?

What are the features of the primordi
present ~10 μs after the Big

What happ **the Universe**
~10

Are th **forces in addition to the known four ?**
Are there additional (microscopic) space dimensions ?
….

New Physics beyond the Standard Model is needed to answer these and other questions. This New Physics could manifest itself at the ~ TeV energy scale being explored by the LHC

How the LHC came to be …

(see a nice article by Chris Llewellyn Smith in Nature 448, p281)

Some early key dates

1977 **The community talked about the LEP project, and it was already mentioned that a new tunnel could also house a hadron collider in the far future**

1981 **LEP was approved with a large and long (27 km) tunnel**

1983 **The early 1980s were crucial:**

The real belief that a 'dirty' hadron collider can actually do great discovery physics came from UA1 and UA2 with their W and Z boson discoveries at CERN



A very early Z → ee online display from one of the detectors (UA2)

1984 For the community it all started with the CERN - ECFA Workshop in Lausanne on the feasibility of a hadron collider in the future LEP tunnel

1987 La Thuile Workshop

Many LHC colleagues were already involved in this WS set up by Carlo Rubbia as part of the Long Range Planning Committee

1989 ECFA Study Week in Barcelona for LHC instrumentation

1990 Large Hadron Collider Workshop Aachen (CERN - ECFA)

1992 CERN – ECFA meeting 'Towards the LHC Experimental Programme' in Evian


P. Jenni

ECFA 84/85
CERN 84–10
5 September 1984

LARGE HADRON COLLIDER
IN THE LEP TUNNEL

Vol. I

PROCEEDINGS OF THE ECFA–CERN WORKSHOP

held at Lausanne and Geneva,
21–27 March 1984

ATLAS and CMS were born with Letters of Intent (LoI), submitted on 1st October 1992, more than 20 years ago

ATLAS and CMS were born with Letters of Intent (LoI), submitted on 1st October 1992, more than 20 years ago

Spokesperson Fabiola Gianotti, celebrating 20 years of ATLAS on 1st October 2012

1991 December CERN Council:
'LHC is the right machine for advance of the subject and the future of CERN' (thanks to the great push by DG C Rubbia)

1993 December proposal of LHC with commissioning in 2002

1994 June Council:

Staged construction was proposed by DG Chris Llewellyn Smith, but some countries could not yet agree, so the Council session vote was suspended until

16 December 1994 Council:

(Two-stage) construction of LHC was approved



The two-stage approval of LHC was understood to be modified in case sufficient CERN non-member state contributions would become available

A lot of LHC campaigns and negotiations took place in the years 1995 - 1997, including also the experiments

Japan, Russia, India, Canada and the USA were agreeing in that phase to contribute to the LHC

(Israel contributed all along to the full CERN programme and LHC)

1997

December Council approved finally the single-stage 14 TeV LHC for completion in 2005

Delivery of the last dipole for the LHC injection lines from Russia (15th June 2001), with L Maiani and A Skrinsky in the centre

The LHC machine



The Large Hadron Collider is a 27 km long collider ring housed in a tunnel about 100 m underground near Geneva

The first cyclotron, and the famous 184” one of Berkeley

Ernest Lawrence
(1901 - 1958)

The first circular accelerator (Berkeley 1930)

184” 1946



The most challenging components were the 1232 high-tech superconducting dipole magnets

Magnetic field:	**8.4 T**
Operation temperature:	**1.9 K**
(120 tons of superfluid Helium)	
Dipole current:	**11700 A**
Stored energy:	**7 MJ**
Dipole weight:	**34 tons**

7600 km of Nb-Ti superconducting cable



$p(\text{TeV}) = 0.3\, B(\text{T})\, R(\text{km})$

LHC Construction Project Leader Lyndon Evans

The LHC is the largest cryogenic system on earth, cooler than outer space



One sector: 3.3 km, 154 dipoles

Magnets cooled down in a bath of ~120 tons of superfluid Helium (excellent thermal conductor)

H K Onnes
Nobel Prize in Physics 1913



- **105 years ago, on 10 July 1908: Heike K Onnes first liquefied Helium (60 ml in 1 hour) in Leiden**
- **LHC today: 32000 He liters liquefied per hour by eight big cryogenic plants (the largest refrigerator in the world)**

The particle beams are accelerated by superconducting Radio-Frequency (RF) cavities





Note: The acceleration is not such a big issue in pp colliders (unlike in e^+e^- colliders), because of the ~ $1/m^4$ behaviour of the synchrotron radiation energy losses [~ E^4_{beam}/Rm^4]

Special quadrupole magnets ('Inner Triplets') are focussing the particle beams to reach highest densities ('luminosity') at their interaction point in the centre of the experiments



Relative beam sizes around the collision point

CERN's particle accelerator chain



Eine am 30. März 2010 aufgezeichnete Teilchenkollision in LHC

Collisions at LHC



Proton-Proton

Protons/bunch	10^{11}
Beam energy	7 TeV (7×10^{12} eV)
Luminosity	10^{34} cm^{-2} s^{-1}

Event rate:

N = L x σ (pp) ≈ 10^9 interactions/s

Mostly soft (low p_T) events

Interesting hard (high-p_T) events are rare

Selection of 1 in 10,000,000,000,000

→ very powerful detectors needed

The SM is not a complete theory

Some of the outstanding questions in fundamental physics are

What is the origin of the elementary particle masses ?	**ATLAS, CMS**
What is the nature of the Universe dark matter ?	**ATLAS, CMS**
Why is only matter observed in the Universe as primary constituents and not anti-matter ?	**LHCb**
What are the features of the primordial plasma present ~10 μs after the Big Bang ?	**ALICE**
What happened in the first moments of the Universe ~10^{-11} s after the Big Bang ?	**ATLAS, CMS**
Are there other forces in addition to the known four ? Are there additional (microscopic) space dimensions ? ….	**ATLAS, CMS**

The LHC World of CERN

Plus smaller
local earldoms
LHCf (point-1)
TOTEM (point-5)
Moedal (point-8)



CMS
3000 Physicists
184 Institutions
38 countries
550 MCHF

LHCb
730 Physicists
54 Institutions
15 countries
75 MCHF

ATLAS
3000 Physicists
177 Institutions
38 countries
550 MCHF

ALICE
1300 Physicists
130 Institutions
35 countries
160 MCHF

Specialized detectors



General purpose detectors



Magnetic fields

Magnetic field configurations:





From C.Joram

Exploded View of CMS



An Example of an Engineering Challenge: CMS Solenoid



CMS solenoid:

Magnetic length	**12.5 m**
Diameter	**6 m**
Magnetic field	**4 T**
Nominal current	**20 kA**
Stored energy	**2.7 GJ**

Tested at full current in Summer 2006



CMS before closure 2008

ATLAS Collaboration

38 Countries
177 Institutions
3000 Scientific participants total
(1000 Students)

Adelaide, Albany, Alberta, NIKHEF Amsterdam, Ankara, LAPP Annecy, Argonne NL, Arizona, UT Arlington, Athens, NTU Athens, Baku, IFAE Barcelona, Belgrade, Bergen, Berkeley LBL and UC, HU Berlin, Bern, Birmingham, UAN Bogota, Bologna, Bonn, Boston, Brandeis, Brasil Cluster, Bratislava/SAS Kosice, Brookhaven NL, Buenos Aires, Bucharest, Cambridge, Carleton, CERN, Chinese Cluster, Chicago, Chile, Clermont-Ferrand, Columbia, NBI Copenhagen, Cosenza, AGH UST Cracow, IFJ PAN Cracow, SMU Dallas, UT Dallas, DESY, Dortmund, TU Dresden, Dubna, Duke, Edinburgh, Frascati, Freiburg, Geneva, Genoa, Giessen, Glasgow, Göttingen, LPSC Grenoble, Technion Haifa, Hampton, Harvard, Heidelberg, Hiroshima IT, Indiana, Innsbruck, Iowa SU, Iowa, UC Irvine, Istanbul Bogazici, KEK, Kobe, Kyoto, Kyoto UE, Kyushu, Lancaster, UN La Plata, Lecce, Lisbon LIP, Liverpool, Ljubljana, QMW London, RHBNC London, UC London, Louisiana Tech, Lund, UA Madrid, Mainz, Manchester, CPPM Marseille, Massachusetts, MIT, Melbourne, Michigan, Michigan SU, Milano, Minsk NAS, Minsk NCPHEP, Montreal, McGill Montreal, RUPHE Morocco, FIAN Moscow, ITEP Moscow, MEPhI Moscow, MSU Moscow, LMU Munich, MPI Munich, Nagasaki IAS, Nagoya, Naples, New Mexico, New York, Nijmegen, Northern Illinois, BINP Novosibirsk, Ohio SU, Okayama, Oklahoma, Oklahoma SU, Olomouc, Oregon, LAL Orsay, Osaka, Oslo, Oxford, Paris VI and VII, Pavia, Pennsylvania, NPI Petersburg, Pisa, Pittsburgh, CAS Prague, CU Prague, TU Prague, IHEP Protvino, Rome I, Rome II, Rome III, Rutherford Appleton Laboratory, DAPNIA Saclay, Santa Cruz UC, Sheffield, Shinshu, Siegen, Simon Fraser Burnaby, SLAC, South Africa, Stockholm, KTH Stockholm, Stony Brook, Sydney, Sussex, AS Taipei, Tbilisi, Tel Aviv, Thessaloniki, Tokyo ICEPP, Tokyo MU, Tokyo Tech, Toronto, TRIUMF, Tsukuba, Tufts, Udine/ICTP, Uppsala, UI Urbana, Valencia, UBC Vancouver, Victoria, Warwick, Waseda, Washington, Weizmann Rehovot, FH Wiener Neustadt, Wisconsin, Wuppertal, Würzburg, Yale, Yerevan

ATLAS Collaboration

38 Countries
177 Institutions
3000 Scientific participants total
(1000 Students)

It is a great pleasure to collaborate with ~425 colleagues from Germany, junior and senior, from 13 universities, DESY and MPI Munich

(GK1504/2 teams in red)

Adelaide, Albany, Alberta, NIKHEF Amsterdam, Ankara, LAPP Annecy, Argonne NL, Arizona, UT Arlington, Athens, NTU Athens, Baku, IFAE Barcelona, Belgrade, Bergen, Berkeley LBL and UC, **HU Berlin**, Bern, Birmingham, UAN Bogota, Bologna, Bonn, Boston, Brandeis, Brasil Cluster, Bratislava/SAS Kosice, Brookhaven NL, Buenos Aires, Bucharest, Cambridge, Carleton, CERN, Chinese Cluster, Chicago, Chile, Clermont-Ferrand, Columbia, NBI Copenhagen, Cosenza, AGH UST Cracow, IFJ PAN Cracow, SMU Dallas, UT Dallas, **DESY**, Dortmund, **TU Dresden**, Dubna, Duke, Edinburgh, Frascati, Freiburg, Geneva, Genoa, Giessen, Glasgow, Göttingen, LPSC Grenoble, Technion Haifa, Hampton, Harvard, Heidelberg, Hiroshima IT, Indiana, Innsbruck, Iowa SU, Iowa, UC Irvine, Istanbul Bogazici, KEK, Kobe, Kyoto, Kyoto UE, Kyushu, Lancaster, UN La Plata, Lecce, Lisbon LIP, Liverpool, Ljubljana, QMW London, RHBNC London, UC London, Louisiana Tech, Lund, UA Madrid, Mainz, Manchester, CPPM Marseille, Massachusetts, MIT, Melbourne, Michigan, Michigan SU, Milano, Minsk NAS, Minsk NCPHEP, Montreal, McGill Montreal, RUPHE Morocco, FIAN Moscow, ITEP Moscow, MEPhI Moscow, MSU Moscow, LMU Munich, MPI Munich, Nagasaki IAS, Nagoya, Naples, New Mexico, New York, Nijmegen, Northern Illinois, BINP Novosibirsk, Ohio SU, Okayama, Oklahoma, Oklahoma SU, Olomouc, Oregon, LAL Orsay, Osaka, Oslo, Oxford, Paris VI and VII, Pavia, Pennsylvania, NPI Petersburg, Pisa, Pittsburgh, CAS Prague, CU Prague, TU Prague, IHEP Protvino, Rome I, Rome II, Rome III, Rutherford Appleton Laboratory, DAPNIA Saclay, Santa Cruz UC, Sheffield, Shinshu, Siegen, Simon Fraser Burnaby, SLAC, South Africa, Stockholm, KTH Stockholm, Stony Brook, Sydney, Sussex, AS Taipei, Tbilisi, Tel Aviv, Thessaloniki, Tokyo ICEPP, Tokyo MU, Tokyo Tech, Toronto, TRIUMF, Tsukuba, Tufts, Udine/ICTP, Uppsala, UI Urbana, Valencia, UBC Vancouver, Victoria, Warwick, Waseda, Washington, Weizmann Rehovot, FH Wiener Neustadt, Wisconsin, Wuppertal, Würzburg, Yale, Yerevan

Age distribution of the ATLAS population





ATLAS

Length : ~ 46 m
Radius : ~ 12 m
Weight : ~ 7000 tons
~ 10^8 electronic channels
~ 3000 km of cables

- **Tracking ($|\eta|<2.5$, B=2T) :**
  - **-- Si pixels and strips**
  - **-- Transition Radiation Detector (e/π separation)**
- **Calorimetry ($|\eta|<5$) :**
  - **-- EM : Pb-LAr**
  - **-- HAD: Fe/scintillator (central), Cu/W-LAr (fwd)**
- **Muon Spectrometer ($|\eta|<2.7$) :**
  - **-- air-core toroids with muon chambers**



The Underground Cavern at Point-1 for the ATLAS Detector



Length = 55 m
Width = 32 m
Height = 35 m



LHC Point 1 - UX 15 Cavern - Concrete walls 6th lift - 20-02-2003 - CERN ST-CE





Hector Berlioz, “Les Troyens”, opera in five acts
Valencia, Palau de les Arts Reina Sofia, 31 October -12 November 2009


ATLAS 16 Feb 2007


ATLAS 4 April 2008
Peter Higgs

Interconnections of two magnets

One (superconductor) joint failed on 19th September 2008, and it caused a catastrophic He-release that made serious collateral damage to sector 3-4 of the LHC machine

The joy in the ATLAS Control Room when the first LHC beam collided on November 23rd, 2009....

First collisions at the LHC end of November 2009 with beams at the injection energy of 450 GeV





A well-deserved toast to all who have built such a marvelous machine, and to all who operate it so superbly

(first 7 TeV collisions on 30th March 2010)

The LHC and experiments performances were simply fantastic over the last three years

Total integrated luminosity

$$N_{events} = \sigma \int L\, dt$$





The experiment records typically 94% of the stably delivered luminosity, and uses up to 90% of the LHC luminosity in the final analyses!

Excellent LHC performance is a (nice) challenge for the experiment:

- **Trigger** *(GK1504 expertize!)*
- **Pile-up**
- **Maintain accuracy of the the measurements in this environment**





Inner Detector for a Z → μμ event with 25 primary vertices

The Worldwide LHC Computing Grid (wLCG)



German Tier-1 at KIT Karlsruhe

DESY Tier-2 with HUB

Tier-0 (CERN):
- Data recording
- Initial data reconstruction
- Data distribution

Tier-1 (12 centres):
- Permanent storage
- Re-processing
- Analysis
- Simulation

Tier-2 (68 federations of >100 centres):
- Simulation
- End-user analysis

Physics Highlights

ATLAS and CMS have already published together about 550 papers in scientific journals (and many more as public conference notes…)

The other experiments, ALICE, LHCb, LHCf, and TOTEM total another 220 journal publications together

It is clearly not possible to cover all these results…

No attempt is made to show in a democratic way, for example, CMS and ATLAS results, only examples are given that are meant to represent the others as well where applicable...

Note that all public results are available from the experiments Web pages, and from the CERN Document Server
http://cdsweb.cern.ch/collection/LHC%20Experiments?ln=en





Physics Highlights:

General event properties

Heavy flavour physics

Standard Model physics including QCD jets

Higgs searches

Searches for SUSY

Searches for 'exotic' new physics



2010

Data corresponding to ~40 pb^{-1} collected → re-discovery of the Standard Model



The di-muon spectrum recalls a long period of particle physics:
Well known quark-antiquark resonances (bound states) appear "online"

2010

Data corre cted
→ re-disco el



The di-muon spectrum recalls a long period of particle physics:
Well known quark-antiquark resonances (bound states) appear "online"

Jet physics


ATLAS
EXPERIMENT
Run Number: 209580, Event Number: 179229707
Date: 2012-08-31 20:24:29 CEST
m_{jj} = 4.7 TeV
p_T^j = 2.3 TeV
E_T^{miss} = 47 GeV
325 ET (GeV)
285
245
205
165
125
85
45
5
η
Φ
0
90
180
270
360

Note also that the event displays have become more sophisticated since the first spectacular events, hand-drawn, at a hadron collider …

Volume 118B, number 1, 2, 3 PHYSICS LETTERS 2 December 1982





ATLAS, EPJC 73 (2013) 2509

Very detailed jet measurements are now available from LHC that can be compared with QCD calculations …

Example: The inclusive jet cross sections as a function of the jet P_T in rapidity bins



CMS-PAS-SMP-12-012

Very detailed jet measurements are now available from LHC that can be compared with QCD calculations …

Example: The inclusive jet cross sections as a function of the jet P_T in rapidity bins



CMS-PAS-SMP-12-012

Cross-section ratios of multi-jets allow one to determine α_s

$$R_{3/2}(p_T^{lead}) = \frac{d\sigma_{N_{jet}\geq 3}/dp_T^{lead}}{d\sigma_{N_{jet}\geq 2}/dp_T^{lead}}$$

$p_T > 40$ GeV and $|y| < 2.8$.



$$\alpha_s(M_Z) = 0.111 \pm 0.006(\text{exp.})\ ^{+0.016}_{-0.003}(\text{theory}).$$



ATLAS-CONF-2013-041

Standard Model Physics

Candidate Z → $\mu^+\mu^-$



W → eν candidate



Today each ATLAS and CMS have in their data more than:

100 M W→ μν, eν events
10 M Z → μμ, ee events

after all selection cuts

Z and W production

Phys Rev D85 (2012) 072004





JHEP 10 (2011) 132

Z peak (di-lepton pair mass distributions, can be extracted essentially background-free)

$$m = \sqrt{(E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2}$$

W transverse mass



μ with p_T>20 GeV, E_T^{miss}>25 GeV



Missing transverse energy from the W → μ + ν decays

ATLAS-CONF-2011-041

$$m_T = \sqrt{2 p_T^{\ell} p_T^{\nu} (1 - \cos(\phi^{\ell} - \phi^{\nu}))}$$

What a contrast to the Intermediate Vector Boson discovery distributions in 1982 and 1983 by UA1 and UA2 …



30 years ago!

(here are shown the UA2 distributions)

Cross section measurements



CMS-PAS-SMP-12-011


$p_T(\mu)$ = 18 GeV
$p_T^{vis}(\tau_h)$ = 26 GeV
$m_{vis}(\mu,\tau_h)$ = 47 GeV
$m_T(\mu,E_T^{miss})$ = 8 GeV
E_T^{miss} = 7 GeV
ATLAS EXPERIMENT
Run Number: 160613, Event Number: 9209492
Date: 2010-08-03 02:12:37 CEST
$Z \rightarrow \tau\tau$
Candidate in 7 TeV Collisions
muon
2 GeV
-5
-3
-1
η
1
3
5
0
180
360
φ
3-prong hadronic tau decay

$p_T(\mu) = 18$ GeV
$p_T^{vis}(\tau_h) = 26$ GeV
$m_{vis}(\mu,\tau_h) = 47$ GeV
$m_T(\mu,E_T^{miss}) = 8$ GeV
$E_T^{miss} = 7$ GeV

ATLAS EXPERIMENT

Run Number: 160613, Event Number: 9209492

Date: 2010-08-03 02:12:37 CEST

$Z \rightarrow \tau\tau$ Candidate in 7 TeV Collisions



muon



GK1504 expertize!



3-prong hadronic tau decay

Top measurements

- **Complete set of ingredients to investigate production of ttbar, which is the next step in verifying the SM at the LHC:**
  - **e, μ, E_T^{miss}, jets, b-tag**

- **Assume all tops decay to Wb: event topology then depends on the W decays:**
  - **one lepton (e or μ), E_T^{miss}, jjbb (37.9%)**
  - **di-lepton (ee, μμ or eμ), E_T^{miss}, bb (6.5%)**



- **Data-driven methods to control QCD and W+jets backgrounds**

$t\bar{t}$ candidate event

e + μ + 2 jets (b-tagged) +ETmiss





$t\bar{t}$ pair production cross-sections

ATLAS-CONF-2012-149
CMS-PAS-TOP-12-006, 007

Top quark mass measurements

ATLAS-CONF-2013-102
CMS-PAS-TOP-13-005



A summary of Standard Model measurements



Similar impressive results from CMS

GK1504 expertize!

The excellent performance in measuring Standard Model physics gives confidence for the readiness of the two experiments to search for New Physics

HIGGS BOSON

A most basic question is why particles (and matter) have masses (and so different masses)

The mass mystery for fundamental particles could be solved with the 'EW symmetry breaking mechanism' which predicts the existence of a new elementary particle, the 'Higgs' particle (theory 1964: R. Brout and F. Englert; P.W. Higgs; G.S. Guralnik, C.R. Hagen and T.W.B. Kibble)

Peter Higgs



The Higgs (H) particle has been searched for since decades at accelerators …

The LHC has sufficient energy to produce it for sure, if it exists

Francois Englert

A cartoon 'illustrating' the scalar Brout-Englert-Higgs field filling all space that affects elementary particles, and 'gives' them their mass by interacting with them



PhD comics © Jorge Cham
http://www.phdcomics.com/higgs/

Search for the boson (H) of the EW symmetry breaking

SM H boson production cross sections times observable decay branching ratios at 8 TeV









Discovery of the Higgs boson



Candidate event for H → ZZ* → ee μμ

Candidate event for H → γγ



ATLAS and CMS have announced the discovery of a new boson together on 4th July 2012, published in a special issue of Physics Letter B



Phys. Lett. B 716 (2012) 1

Phys. Lett. B 716 (2012) 30

Very happy faces after the announcement of the discovery on 4th July 2012 at CERN and at ICHEP Melbourne

ICHEP2012 Melbourne

H → γγ

- Small cross-section: σ ~ 40 fb
- Expected S/B ~ 0.02
- Simple final state: two high-p_T isolated photons
- Main background: γγ continuum (irreducible) and fake γ from γj and jj events (reducible)



ATLAS-CONF-2013-012 and Phys. Lett. B 726 (2013) 88-119





CMS-PAS-HIG-13-001

$H \rightarrow ZZ^{(*)} \rightarrow 4l$ (4e, 4μ, 2e2μ)

- Rare process, small cross section: σ ~ 2-5 fb
- However: pure: S/B ~ 1
- 4 leptons:
- Main background: $ZZ^{(*)}$ (irreducible)

  In addition: Zbb, Z+jets, tt with two leptons from b-quarks or jets



ATLAS-CONF-2013-013 and Phys. Lett. B 726 (2013) 88-119



CMS-PAS-HIG-13-002

H → WW$^{(*)}$ → lνlν (eνeν, μνμν, eνμν)

- Very sensitive channel over ~ 125-180 GeV (σ ~ 200 fb)
- Challenging: 2ν → no mass reconstruction/peak → "counting channel"
- 2 isolated opposite-sign leptons, use eνμν only for 2012 data, large E_T^{miss}
- Main backgrounds: WW, top, Z+jets, W+jets
- Topological cuts against "irreducible" WW background

(Just an example distributions from several categories used in both experiments)



ATLAS-CONF-2013-030 and Phys. Lett. B 726 (2013) 88-119

$$m_T = \sqrt{(E_T^{\ell\ell} + E_T^{miss})^2 - (\mathbf{P}_T^{\ell\ell} + \mathbf{P}_T^{miss})^2}$$

How significant is the signal for the new particle ?

Mass = 125.5 ± 0.2 (stat) ± 0.6 (syst) GeV [ATLAS]
125.7 ± 0.3 (stat) ± 0.3 (syst) GeV [CMS]

Observed data compared to the probability that the background fluctuates to fake the observed excess of events, and what is expected from a SM Higgs

Signal strength
$\mu = 0$ background only hypothesis
$\mu = 1$ SM Higgs hypothesis





ATLAS-CONF-2013-034

CMS-PAS-HIG-13-005

$\mu = 1.30 \pm 0.20$ [ATLAS]
$\mu = 0.80 \pm 0.14$ [CMS]

How significant is the signal for the new particle ?

Mass = 125.5 ± 0.2 (stat) ± 0.6 (syst) GeV [ATLAS]
125.7 ± 0.3 (stat) ± 0.3 (syst) GeV [CMS]

Mass measurements in the two high-resolution channels from CMS



ATLAS-CONF-2013-034

CMS-PAS-HIG-13-005

Signal strength

μ = 0 background only hypothesis

μ = 1 SM Higgs hypothesis



μ = 1.30 ± 0.20 [ATLAS]

μ = 0.80 ± 0.14 [CMS]

Is the new particle a Higgs boson ?

ATLAS and CMS have verified the two "fingerprints"

1) To accomplish its job (providing mass) it interacts with other particles (in particular W, Z) with strength proportional to their masses



2) It has spin zero (scalar)



L ($J^P=0^+$)/L($J^P=2^+$)

Hypothesis	Rejection (C.L.)
0^-	97.8%
1^+	99.97%
1^-	99.7%
2^+	99.9%

From F Gianotti

Detailed studies of the production and decay properties have started in order to characterize the new particle

It will be important to understand with great precision if it is the only scalar boson of the Standard Model 'Brout-Englert-Higgs' mechanism to break the electroweak symmetry, or if it is only part of a broader physics picture going *Beyond the Standard Model*

These studies will be among the most central ones in the decades to come both at the LHC and at possible other future colliders

For the experts:

Couplings
Production modes
Spin-parity

all support at the 2-3 σ level the SM Higgs with present limited statistics

Birth and evolution of a signal: H→4l



Searches Beyond the Standard Model
(only very few examples out of many…)



N C Flammarion 1888
(colours added later)

Supersymmetry (SUSY)

(Julius Wess and Bruno Zumino, 1974)

Establishes a symmetry between fermions (matter) and bosons (forces):

- Each particle p with spin s has a SUSY partner $\tilde{p}$ with spin s -1/2

- Examples **q (s=1/2) → $\tilde{q}$ (s=0) squark**
g (s=1) → $\tilde{g}$ (s=1/2) gluino



Our known world… **Maybe a new world?**



Motivation:

- Unification (fermions-bosons, matter-forces)
- Solves some deep problems of the Standard Model

Dark Matter in the Universe

Astronomers found that most of the matter in the Universe must be invisible Dark Matter

Vera Rubin ~ 1970

'Supersymmetric' particles ?

F. Zwicky 1898-1974



In practice SUSY searches at LHC are rather complicated

- **Complex (and model-dependent) squark/gluino cascades**



Missing Transverse Energy



- **Focus on signatures covering large classes of models while strongly rejecting SM background**
  - **large missing E_T**
  - **High transverse momentum jets**
  - **Leptons**
    - **Perform separate analyses with and without lepton veto (0-lepton / 1-lepton / 2-leptons)**
  - **B-jets: to enhance sensitivity to third-generation squarks**
  - **Photons: typically for models with the gravitino as LSP**

Meff = Etmiss + Σ pT(jets)

An <u>example</u> from the 2011 data, to show the principle, final results will be quoted for updated analyses including 2012 data

- 0-lepton + 2−6 jets + high MET (based on Et-miss+jet triggers)
- 0-lepton + 6−9 (multi-)jets + MET (based on multi-jet triggers)
- 1-lepton + 3,4 jets + high MET (based on lepton triggers)

Example: O-leptons + 2-6 Jets analysis



A signal region

ATLAS-CONF-2012-033, 037, and 041



A control region where no signal is expected

Interpretation of the results

Consider phenomenological MSSM models containing only squarks of 1st and 2nd generation, gluino and light neutralinos



ATLAS-CONF-2013-047

Expected production cross-sections at LHC



ATLAS SUSY Searches* - 95% CL Lower Limits

Status: LP 2013

SUSY limits

ATLAS Preliminary

$\int \mathcal{L}\, dt = (4.4 - 22.9)\ \mathrm{fb}^{-1}$ $\sqrt{s} = 7, 8$ TeV

	Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L}\, dt[\mathrm{fb}^{-1}]$	Mass limit		Reference
Inclusive Searches	MSUGRA/CMSSM	1 e, μ	3-6 jets	Yes	20.3	$\tilde{g}$ 1.2 TeV	any $m(\tilde{q})$	ATLAS-CONF-2013-062
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	$\tilde{g}$ 1.1 TeV	any $m(\tilde{q})$	ATLAS-CONF-2013-054
	$\tilde{q}\tilde{q}, \tilde{q}\to q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{q}$ 740 GeV	$m(\tilde{\chi}_1^0)$=0 GeV	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g}\to q\bar{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{g}$ 1.3 TeV	$m(\tilde{\chi}_1^0)$=0 GeV	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g}\to qq\tilde{\chi}_1^\pm\to qqW^\pm\tilde{\chi}_1^0$	1 e, μ	3-6 jets	Yes	20.3	$\tilde{g}$ 1.18 TeV	$m(\tilde{\chi}_1^0)$<200 GeV, $m(\tilde{\chi}^\pm)=0.5(m(\tilde{\chi}_1^0)+m(\tilde{g}))$	ATLAS-CONF-2013-062
	$\tilde{g}\tilde{g}\to qqqq\ell\ell(\ell\ell)\tilde{\chi}_1^0\tilde{\chi}_1^0$	2 e, μ (SS)	3 jets	Yes	20.7	$\tilde{g}$ 1.1 TeV	$m(\tilde{\chi}_1^0)$<650 GeV	ATLAS-CONF-2013-007
	GMSB ($\tilde{\ell}$ NLSP)	2 e, μ	2-4 jets	Yes	4.7	$\tilde{g}$ 1.24 TeV	$\tan\beta$<15	1208.4688
	GMSB ($\tilde{\ell}$ NLSP)	1-2 τ	0-2 jets	Yes	20.7	$\tilde{g}$ 1.4 TeV	$\tan\beta$ >18	ATLAS-CONF-2013-026
	GGM (bino NLSP)	2 γ	0	Yes	4.8	$\tilde{g}$ 1.07 TeV	$m(\tilde{\chi}_1^0)$>50 GeV	1209.0753
	GGM (wino NLSP)	1 $e, \mu + \gamma$	0	Yes	4.8	$\tilde{g}$ 619 GeV	$m(\tilde{\chi}_1^0)$>50 GeV	ATLAS-CONF-2012-144
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	4.8	$\tilde{g}$ 900 GeV	$m(\tilde{\chi}_1^0)$>220 GeV	1211.1167
	GGM (higgsino NLSP)	2 e, μ (Z)	0-3 jets	Yes	5.8	$\tilde{g}$ 690 GeV	$m(\tilde{H})$>200 GeV	ATLAS-CONF-2012-152
	Gravitino LSP	0	mono-jet	Yes	10.5	$F^{1/2}$ scale 645 GeV	$m(\tilde{g})>10^{-4}$ eV	ATLAS-CONF-2012-147
3rd gen. $\tilde{g}$ med.	$\tilde{g}\to b\bar{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	$\tilde{g}$ 1.2 TeV	$m(\tilde{\chi}_1^0)$<600 GeV	ATLAS-CONF-2013-061
	$\tilde{g}\to t\bar{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	$\tilde{g}$ 1.14 TeV	$m(\tilde{\chi}_1^0)$ [illegible] eV	ATLAS-CONF-2013-054
	$\tilde{g}\to t\bar{t}\tilde{\chi}_1^0$	0-1 e, μ	3 b	Yes	20.1	$\tilde{g}$ 1.34 TeV	[illegible]	ATLAS-CONF-2013-061
	$\tilde{g}\to b\bar{t}\tilde{\chi}_1^+$	0-1 e, μ	3 b	Yes	20.1	$\tilde{g}$ 1.3 TeV	[illegible]	ATLAS-CONF-2013-061
3rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1\to b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	$\tilde{b}_1$ 100-630 GeV		ATLAS-CONF-2013-053
	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1\to t\tilde{\chi}_1^\pm$	2 e, μ (SS)	0-3 b	Yes	20.7	$\tilde{b}_1$ 430 GeV	[illegible] $m(\tilde{\chi}_1^0)$	ATLAS-CONF-2013-007
	$\tilde{t}_1\tilde{t}_1$(light), $\tilde{t}_1\to b\tilde{\chi}_1^\pm$	1-2 e, μ	1-2 b	Yes	4.7	$\tilde{t}_1$ 167 GeV	$m(\tilde{\chi}_1^0)$=55 GeV	1208.4305, 1209.2102
	$\tilde{t}_1\tilde{t}_1$(light), $\tilde{t}_1\to Wb\tilde{\chi}_1^0$	2 e, μ	0-2 jets	Yes	20.3	$\tilde{t}_1$ 220 GeV	$m(\tilde{\chi}_1^0)=m(\tilde{t}_1)-m(W)$-50 GeV, $m(\tilde{t}_1)<<m(\tilde{\chi}_1^\pm)$	ATLAS-CONF-2013-048
	$\tilde{t}_1\tilde{t}_1$(medium), $\tilde{t}_1\to b\tilde{\chi}_1^\pm$	2 e, μ	0-2 jets	Yes	20.3	$\tilde{t}_1$ [illegible]	$m(\tilde{\chi}_1^0)$=0 GeV, $m(\tilde{t}_1)-m(\tilde{\chi}_1^\pm)$=10 GeV	ATLAS-CONF-2013-048
	$\tilde{t}_1\tilde{t}_1$(medium), $\tilde{t}_1\to b\tilde{\chi}_1^\pm$	0	2 b	Yes	20.1	$\tilde{t}_1$ [illegible]	$m(\tilde{\chi}_1^0)$<200 GeV, $m(\tilde{\chi}_1^\pm)-m(\tilde{\chi}_1^0)$=5 GeV	ATLAS-CONF-2013-053
	$\tilde{t}_1\tilde{t}_1$(heavy), $\tilde{t}_1\to t\tilde{\chi}_1^0$	1 e, μ	1 b	Yes	20.7	$\tilde{t}_1$ [illegible] 0 GeV	$m(\tilde{\chi}_1^0)$=0 GeV	ATLAS-CONF-2013-037
	$\tilde{t}_1\tilde{t}_1$(heavy), $\tilde{t}_1\to t\tilde{\chi}_1^0$	0	2 b	Yes	20.5	$\tilde{t}_1$ 320-660 GeV	$m(\tilde{\chi}_1^0)$=0 GeV	ATLAS-CONF-2013-024
	$\tilde{t}_1\tilde{t}_1$(natural GMSB)	2 e, μ (Z)	1 b	Yes	20.7	[illegible] 500 GeV	$m(\tilde{\chi}_1^0)$>150 GeV	ATLAS-CONF-2013-025
	$\tilde{t}_2\tilde{t}_2, \tilde{t}_2\to\tilde{t}_1 + Z$	3 e, μ (Z)	1 b	Yes	[illegible]	[illegible] 520 GeV	$m(\tilde{t}_1)=m(\tilde{\chi}_1^0)$+180 GeV	ATLAS-CONF-2013-025
EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell}\to\ell\tilde{\chi}_1^0$	2 e, μ	0	[illegible]	[illegible]	[illegible] 85-315 GeV	$m(\tilde{\chi}_1^0)$=0 GeV	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+\to\tilde{\ell}\nu(\ell\tilde{\nu})$	2 e, μ	[illegible]	[illegible]	[illegible]	[illegible] 125-450 GeV	$m(\tilde{\chi}_1^0)$=0 GeV, $m(\tilde{\ell}, \tilde{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+\to\tilde{\tau}\nu(\tau\tilde{\nu})$	2 τ	[illegible]	[illegible]	[illegible]	$\tilde{\chi}_1^\pm$ 180-330 GeV	$m(\tilde{\chi}_1^0)$=0 GeV, $m(\tilde{\tau}, \tilde{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-028
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0\to\tilde{\ell}_L\nu\tilde{\ell}_L\ell(\tilde{\nu}\nu), \ell\tilde{\nu}\tilde{\ell}_L\ell(\tilde{\nu}\nu)$	3 e, μ	[illegible]	[illegible]	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 600 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0)$, $m(\tilde{\chi}_1^0)$=0, $m(\tilde{\ell}, \tilde{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0\to W^*\tilde{\chi}_1^0 Z^*\tilde{\chi}_1^0$	3 e, μ	[illegible]	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 315 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0)$, $m(\tilde{\chi}_1^0)$=0, sleptons decoupled	ATLAS-CONF-2013-035
Long-lived particles	Direct $\tilde{\chi}_1^+\tilde{\chi}_1^-$ prod., long-lived $\tilde{\chi}_1^\pm$	0	1 jet	Yes	4.7	$\tilde{\chi}_1^\pm$ 220 GeV	$1<\tau(\tilde{\chi}_1^\pm)<10$ ns	1210.2852
	Stable, stopped $\tilde{g}$ R-hadron	0	1-5 jets	Yes	22.9	$\tilde{g}$ 857 GeV	$m(\tilde{\chi}_1^0)$=100 GeV, 10 μs<$\tau(\tilde{g})$<1000 s	ATLAS-CONF-2013-057
	GMSB, stable $\tilde{\tau}$	1-2 μ	0	-	15.9	$\tilde{\tau}$ 385 GeV	5<$\tan\beta$<50	ATLAS-CONF-2013-058
	Direct $\tilde{\tau}\tilde{\tau}$ prod., stable $\tilde{\tau}$ or $\tilde{\ell}$	1-2 μ	0	-	15.9	$\tilde{\tau}$ 395 GeV	$m(\tilde{\tau})=m(\tilde{\ell})$	ATLAS-CONF-2013-058
	GMSB, $\tilde{\chi}_1^0\to\gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$	2 γ	0	Yes	4.7	$\tilde{\chi}_1^0$ 230 GeV	$0.4<\tau(\tilde{\chi}_1^0)<2$ ns	1304.6310
	$\tilde{\chi}_1^0\to qq\mu$ (RPV)	1 μ	0	Yes	4.4	$\tilde{q}$ 700 GeV	1 mm<$c\tau$<1 m, $\tilde{g}$ decoupled	1210.7451
RPV	LFV $pp\to\tilde{\nu}_\tau + X, \tilde{\nu}_\tau\to e + \mu$	2 e, μ	0	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV	λ'_{311}=0.10, λ_{132}=0.05	1212.1272
	LFV $pp\to\tilde{\nu}_\tau + X, \tilde{\nu}_\tau\to e(\mu) + \tau$	1 $e, \mu + \tau$	0	-	4.6	$\tilde{\nu}_\tau$ 1.1 TeV	λ'_{311}=0.10, $\lambda_{1(2)33}$=0.05	1212.1272
	Bilinear RPV CMSSM	1 e, μ	7 jets	Yes	4.7	$\tilde{q}, \tilde{g}$ 1.2 TeV	$m(\tilde{q})=m(\tilde{g})$, $c\tau_{LSP}$<1 mm	ATLAS-CONF-2012-140
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+\to W\tilde{\chi}_1^0, \tilde{\chi}_1^0\to ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	4 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm$ 760 GeV	$m(\tilde{\chi}_1^0)$>300 GeV, λ_{121}>0	ATLAS-CONF-2013-036
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+\to W\tilde{\chi}_1^0, \tilde{\chi}_1^0\to\tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	3 $e, \mu + \tau$	0	Yes	20.7	$\tilde{\chi}_1^\pm$ 350 GeV	$m(\tilde{\chi}_1^0)$>80 GeV, λ_{133}>0	ATLAS-CONF-2013-036
	$\tilde{g}\to qqq$	0	6 jets	-	4.6	$\tilde{g}$ 666 GeV		1210.4813
	$\tilde{g}\to\tilde{t}_1 t, \tilde{t}_1\to bs$	2 e, μ (SS)	0-3 b	Yes	20.7	$\tilde{g}$ 880 GeV		ATLAS-CONF-2013-007
Other	Scalar gluon	0	4 jets	-	4.6	sgluon 100-287 GeV	incl. limit from 1110.2693	1210.4826
	WIMP interaction (D5, Dirac χ)	0	mono-jet	Yes	10.5	M^* scale 704 GeV	$m(\chi)$<80 GeV, limit of<687 GeV for D8	ATLAS-CONF-2012-147

10^{-1} 1 **Mass scale [TeV]**

$\sqrt{s}$ = 7 TeV full data | $\sqrt{s}$ = 8 TeV partial data | $\sqrt{s}$ = 8 TeV full data

Very similar limits come from CMS

*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1σ theoretical signal cross section uncertainty.

Searches for heavy W and Z like particles

These searches are quite straight-forward, following basically the same analyses as for the familiar W and Z bosons

Z': Di-lepton pairs



ATLAS-CONF-2013-017

W': Lepton + ETmiss



CMS-EXO-12-060

The highest mass di-lepton events from ATLAS



m (e^+e^-) = 1.54 TeV



m ($\mu^+\mu^-$) = 1.84 TeV

Lower mass limits, at 95% CL, for spin-2 Randall-Sundrum Gravitons



ATLAS-CONF-2013-017

R Sundrum
L Randall
F Gianotti

New particles decaying into two photons

Example for a search of extra dimension signals (Kaluza-Klein Graviton in the Randall-Sundrum and Arkani-Hamed, Dimopoulos and Dvali models)







New J Phys 15 (2013) 043007

If theories with Extra-dimensions are true, microscopic black holes could be abundantly produced and observed at the LHC



Simulation of a black hole event with M_{BH} ~ 8 TeV in ATLAS

They decay immediately through Stephen Hawking radiation

If theories with Extra-dimensions are true, microscopic black holes could be abundantly produced and observed at the LHC



CMS Experiment at LHC, CERN
Data recorded: Mon May 23 21:46:26 2011 EDT
Run/Event: 165567 / 347495624
Lumi section: 280
Orbit/Crossing: 73255853 / 3161

A real 'candidate' event of a 'black hole' in CMS with 9 jets and ST = 2.6 TeV

They decay immediately through Stephen Hawking radiation

Search for Microscopic Black Hole production in models wth large extra dimensions

(Arkani-Hamed, Dimopoulos, Dvali)



Decay into many objects (jets, leptons, photons)

Phys Rev D88 (2013) 072001

<u>Examples:</u> (ATLAS) two same sign muons and large multiplicity, (CMS) any three objects

($S_T = \Sigma P_T$: scalar sum of the E_T of the N objects in the event)

No deviation is seen for events with at least 3 objects with > 50 GeV pT



Submitted to JHEP
arXiv:1303.5338v1[hep-ex]

Similar results exist from ATLAS



LHC Schedule and Upgrade



German groups with DESY will play major roles in the high-luminosity upgrade of the experiments





The High Energy Frontier





©Ralph A. Clevenger/CORBIS

Collider options for the high energy frontier

pp colliders

	Years	E_{cm} TeV	Luminosity $10^{34}cm^{-2}s^{-1}$	Int. Luminosity fb^{-1}
Design LHC	2014-21	14	1-2	300
HL-LHC	2024-30	14	5	3000
HE-LHC	>2035	26-33*	2	100-300/y
V-LHC**	>2035	42-100		

* 16-20 T dipole field
** 80 km Tunnel

e+e- colliders

	Years	E_{cm} GeV	Luminosity $10^{34}cm^{-2}s^{-1}$	Tunnel length km
ILC 250	<2030	250	0.75	
ILC 500		500	1.8	~30
ILC 1000		1000		~50
CLIC 500	>2030	500	2.3(1.3)	~13
CLIC 1400		1400(1500)	3.2(3.7)	~27
CLIC 3000		3000	5.9	~48
LEP3	>2024	240	1	LEP/LHC ring
TLEP	>2030	240	5	80 (ring)
TLEP		350	0.65	80 (ring)

Other options:

$\mu^+\mu^-$ and $\gamma\gamma$ colliders with similar physics as e^+e^- colliders

LHeC for ep collisions

See European Strategy Briefing Book for references

Options for preliminary studies at CERN



The journey into new physics territory has just only begun, and for sure, exciting times are ahead of us!



Thank you for your attention

J Blaeu 1664

Further reading:

The Higgs Boson

ARTICLE

Journey in the Search for the Higgs Boson: The ATLAS and CMS Experiments at the Large Hadron Collider

M. Della Negra,[1] P. Jenni,[2] T. S. Virdee[1]*

The search for the standard model Higgs boson at the Large Hadron Collider (LHC) started more than two decades ago. Much innovation was required and diverse challenges had to be overcome during the conception and construction of the LHC and its experiments. The ATLAS and CMS Collaboration experiments at the LHC have discovered a heavy boson that could complete the standard model of particle physics.

Journey in the Search for the Higgs Boson: The ATLAS and CMS Experiments at the Large Hadron Collider
M. Della Negra *et al.*
Science **338**, 1560 (2012);
DOI: 10.1126/science.1230827

http://www.sciencemag.org/content/338/6114/1560.full.html

Spares

LHC Accelerator Challenge: Dipole Magnets





Magnetic Field for Dipoles
p (TeV) = 0.3 B(T) x R(km)

For p = 7 TeV and R = 4.3 km
⇨ B = 8.4 T
⇨ Current 12 kA

Coldest Ring in the Universe ?
1.9 K (CMBR is about 2.7 K)

LHC magnets are cooled with pressurized superfluid Helium



The LHCb detector installed at Point-8



ALICE (January 2008)



A historical moment:
Closure of the LHC beam pipe ring on 16th June 2008

LHC and ATLAS have also been operated very successfully as Pb-Pb and as p-Pb colliders





Subdetector	Number of Channels	Approximate Operational Fraction
Pixels	80 M	95.9%
SCT Silicon Str		
TRT Transition		
LAr EM Calorin		
Tile calorimeter		
Hadronic endca		
Forward LAr ca		
LVL1 Calo trigg		
LVL1 Muon RP		
LVL1 Muon TG		
MDT Muon Drif		
CSC Cathode S		
RPC Barrel Mu		
TGC Endcap M		



Worldwide LHC Computing Grid (wLCG)



WLCG is a worldwide collaborative effort on an unprecedented scale in terms of storage and CPU requirements, as well as the software project's size



GRID computing developed to solve problem of data storage and analysis

LHC data volume per year: 10-15 Petabytes

One CD has ~ 600 Megabytes
1 Petabyte = 10^9 MB = 10^{15} Byte

(Note: the WWW is from CERN...)

The high quality of the WLCG computing system allows LHC experiments to show results from data taken just few weeks before



Data preparation (ATLAS):

- **First-pass reconstruction at Tier-0 within ~2 days**
- **Calibration good for physics analysis on Grid within ~1 week**





Physics Highlights:

General event properties

Heavy flavour physics

Standard Model physics including QCD jets

Higgs searches

Searches for SUSY

Searches for 'exotic' new physics



Total cross-section measurement by TOTEM





Presented at HCP2012

The first new particles 'discovered' at LHC, December 2011

$\chi_b(3P) \rightarrow \Upsilon(1s,2s)\,\gamma$

m [$\chi_b(3P)$] =10.530 ± 0.005 (stat) ± 0.009 (syst) GeV



Phys. Rev. Lett. 108 (2012) 152001



- $\chi_b(nP) \rightarrow \Upsilon(1s,2s)\,\gamma \rightarrow \mu\mu\,\gamma$
- $X_b(1P)$ m= 9.9 GeV and $X_b(2P)$ m= 10.2 GeV states clearly visible
- New structure at 10.5 GeV → $X_b(3P)$
- Significance > 6 σ
- As theoretically predicted

Two examples of confronting the 2010 data with SM theory





Phys Rev D85 (2012) 072004

Lepton charge asymmetry from W decays in pp collisions at 7 TeV

$$\mathcal{A}(\eta) = \frac{d\sigma/d\eta(W^+ \to \ell^+\nu) - d\sigma/d\eta(W^- \to \ell^-\bar{\nu})}{d\sigma/d\eta(W^+ \to \ell^+\nu) + d\sigma/d\eta(W^- \to \ell^-\bar{\nu})},$$



ATLAS-CONF-2011-129
LHCb-CONF-2011-039
CMS-EWK-10-006 (aXiv:1103.3407)

W → τν signal



CMS-PAS-EWK-11-019

tt – production with b-tagged e-μ events



ATLAS-CONF-2013-097


ATLAS
EXPERIMENT
Run 166466 Event 26227945
Time 2010-10-07 22:16:39 UTC
μ⁻
e
MET
μ⁺
WZ→eνμμ Candidate

Higgs production cross-sections at 8 TeV, and branching fractions



LHC Higgs cross-section working group, arXiv: 1101.0593 and 1201.3084
(the theoretical uncertainties are indicated by the width of the curves)

$H \rightarrow ZZ^{(*)} \rightarrow 4l$ (4e, 4μ, 2e2μ)

- Rare process, small cross section: σ ~ 2-5 fb
- However: pure: S/B ~ 1
- 4 leptons:
- Main background: $ZZ^{(*)}$ (irreducible)
  In addition: Zbb, Z+jets, tt with two leptons from b-quarks or jets

Why a Z peak ?





ATLAS-CONF-2013-013 and Phys. Lett. B 726 (2013) 88-119



CMS-PAS-HIG-13-002

Birth and evolution of a signal: H→ γγ



Two of the by now historical plots from the July 2012 discovery announcement





Observed data compared to the probability that the background fluctuates to fake the observed excess of events, and what is expected from a SM Higgs



A dream becoming true much faster than anticipated long ago



H search in the bb channel





From F Gianotti

Mass measurement

From high-resolution
H→ γγ and H→ 4l channels

$$m_H(\text{combined}) = 125.5\ \text{GeV} \pm 0.2\ (\text{stat})\ ^{+0.5}_{-0.6}(\text{syst})\ \text{GeV}$$



Signal production strength

μ = measured signal production rate in a given final state normalized to SM Higgs expectation

From di-boson final states (γγ, 4l, WW) :
μ = 1.33 ± 0.20

Including preliminary results from fermionic final states (bb, ττ):
μ = 1.23 ± 0.18

→in agreement with SM expectation

Note: similar contributions from statistical and systematic uncertainties to total error and from theory and experimental uncertainties to total systematic error

Constraining production modes and couplings (examples …)

From F Gianotti





$$k_i^2 = \frac{\Gamma_i^{data}}{\Gamma_i^{SM}}$$

Ratios of vector-boson (VBH, VH) to top-quark (ggF, ttH) induced processes



New particles in the gg → H and H→ γγ loops ?



- ❑ > 3σ significance for non-vanishing VBF
- ❑ new particle couples to W and Z as expected
- → first "fingerprint" of a Higgs boson (to do its job → EWSB/Higgs mechanism)
- ❑ No significant New Physics contributions (within present uncertainty)





$$k_i^2 = \frac{\Gamma_i^{data}}{\Gamma_i^{SM}}$$



… and many others …

Main conclusions:

- measurements at the 10-20% level
- couplings to W, Z as expected from EWSB mechanism
- couplings to fermions at > 5σ observed indirectly (mainly through ggF loop)
- no evidence for new physics within present uncertainty

From F Gianotti

From F Gianotti

2nd "fingerprint" of a Higgs boson: spin zero

- Spin information from angular and kinematic distributions of decay products
- H→ γγ, H→ 4l and H→ lνlν channels combined
- Fits to data test $J^P=0^+$ hypothesis (SM) against $J^P= 0^-, 1^+, 1^-, 2^+$

H→ γγ: distribution of polar angle θ* of γγ system for events in signal peak region



Data (after bckg subtraction) **vs**
$J^P=0^+$ (left)
$J^P=2^+$ (right)
expectations



Hypothesis	Rejection (C.L.)
0^-	97.8%
1^+	99.97%
1^-	99.7%
2^+	99.9%

First elementary (likely) scalar observed
→ consequences also for Universe evolution (inflation triggered by a scalar field)

Detailed studies of the production and decay properties have started in order to characterize the new particle

It will be important to understand with great precision if it is the only scalar boson of the Standard Model 'Brout-Englert-Higgs' mechanism to break the electroweak symmetry, or if it is only part of a broader physics picture going *Beyond the Standard Model*

These studies will be among the most central ones in the decades to come both at the LHC and at possible other future colliders

For the experts:

Couplings
Production modes
Spin-parity

all support at the 2-3 σ level the SM Higgs with present limited statistics



ATLAS, Phys. Lett. B 726 (2013) 88-119 and 120-144



CMS-PAS-HIG-13-005

Recall the Happy End of Higgs-Chapter-1 from last month …







"for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at CERN's Large Hadron Collider"

Outlook for HL-LHC on the Higgs physics (I)





ATLAS NOTE

ATL-PHYS-PUB-2012-004

October 15, 2012

Outlook for HL-LHC on the Higgs physics (II)



ttH with H → γγ for 3000 fb^{-1}

H → μμ for 3000 fb^{-1}



ATLAS NOTE

ATL-PHYS-PUB-2012-004

October 15, 2012

Dark Matter in the Universe



Vera Rubin ~ 1970

metric' particles ?

F. Zwicky 1898-1974




$\tilde{g}$-$\tilde{g}$ production, $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, $\sqrt{s}$ = 8 TeV
Status: LHCP 2013
95% CL limits. σ_{theory}^{SUSY} not included.
0-lepton, 7 - ≥ 10 jets [L_{int} = 20.3 fb^{-1}]
ATLAS-CONF-2013-054
0-lepton, ≥ 3 b-jets [L_{int} = 12.8 fb^{-1}]
ATLAS-CONF-2012-145
3-leptons, ≥ 4 jets [L_{int} = 12.8 fb^{-1}]
ATLAS-CONF-2012-151
2-SS-leptons, 0 - ≥ 3 b-jets [L_{int} = 20.7 fb^{-1}]
ATLAS-CONF-2013-007
Expected
Observed
Expected
Observed
Expected
Observed
Expected
Observed
$\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ forbidden
ATLAS Preliminary
$m_{\tilde{\chi}_1^0}$ [GeV]
$m_{\tilde{g}}$ [GeV]
200
400
600
800
1000
500
600
700
800
900
1000
1100
1200
1300

Summary of dedicated searches for top squark pair production for some theoretically preferred models with relatively light 3rd generation squarks



Lower mass limits, at 95% CL, for various Z' and W' like objects



ATLAS-CONF-2013-017



CMS-EXO-12-060

Example of searches for New Physics as deviations from QCD behaviour of hadronic jet distributions

Search for resonances in the di-jet mass spectrum



ATLAS-CONF-2012-148



CMS-EXO-12-059

Example of searches for New Physics as deviations from QCD behaviour of hadronic jet distributions

Search for resonances in the di-jet mass spectrum



ATLAS-CONF-2012-148



CMS-EXO-12-059

Search for direct Dark Matter (DM) particles in pair-production



A single photon (150 GeV) or jet plus ETmiss







ATLAS-CONF-2012-085
arXiv:1210.4491v1[hep-ex]

CMS: Sub. to Phys. Rev. Lett.
arXiv:1204.0821v1[hep-ex]
arXiv:1206.5663[hep-ex]

ATLAS 95% CL limits



*Only a selection of the available mass limits on new states or phenomena shown

Indirect indications for physics BSM, like SUSY, could come from rare decays showing rates deviating from the SM expectations



The search for $B_{s\,(d)} \rightarrow \mu\,\mu$

Submitted to Phys. Rev. Lett.
arXiv:1211.2674v1[hep-ex]

Very rare decay sensitive to New Physics (in particular to models with high *tan β*)

Precise predictions in SM:
$BR(B_s \rightarrow \mu\,\mu) = 3.5 \pm 0.2\ 10^{-9}$
$BR(B_d \rightarrow \mu\,\mu) = 1.1 \pm 0.2\ 10^{-10}$

Very clean experimental signature





With 2011+2012 data (2.1/fb) first evidence of $B_s \rightarrow \mu\,\mu$ decay at ~ 3.5 σ

$\mathcal{B}(B^0_s \rightarrow \mu^+\mu^-) = (3.2^{+1.5}_{-1.2}) \times 10^{-9}$

in agreement with SM. Potential impact on models

Also best limit on $B_d \rightarrow \mu\,\mu$

$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-) < 9.4 \times 10^{-10}$ at 95% CL

2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	...	2030
	Phase 0			LS1		Phase I,II			LS2	Phase II			LS3			

Consolidation	Upgrades

ATLAS has devised a 3 stages program to optimise the physics reach at each Phase

- New Insertable pixel b-layer (IBL)
- New Al beam pipe
- New pixel services
- New evaporative cooling plant
- Consolidation of detector elements (e.g. calorimeter power supplies)
- Add specific neutron shielding
- Finish installation of EE muon chambers staged in 2003
- Upgrade magnet cryogenics

- New Small Wheel (nSW) for the forward muon Spectrometer
- High Precision Calorimeter Trigger at Level-1
- Fast TracKing (FTK) for the Level-2 trigger
- Topological Level-1 trigger processors
- New forward diffractive physics detectors (AFP)

- All new Tracking Detector
- Calorimeter electronics upgrades
- Upgrade part of the muon system
- Possible Level-1 track trigger
- Possible changes to the forward calorimeters