
Fundamentals of Optical Sciences

WS 2015/2016

5 . Exercise

16.11.2015


Lecture: Prof. Dr. Alejandro Saenz, Prof. Dr. Oliver Benson

Prepare your answers for the exercise on 23.11.2015.

Exercise 1

Reflection single dielectric thin film

Light of wavelength λ_0 is incident from air ($n = 1$) onto a single dielectric thin film (of index n_f , and thickness $\lambda/4$, where λ is the wavelength inside the film), which covers a glass substrate (index n_g).



- Write down an expression for the film reflectance, assuming the light is at normal incidence, using the results of the reflection-summation formalism.
- Derive the value of n_f that makes a perfect anti-reflection coating.




Figure 1

Exercise 2

Show that the Liénard-Wiechert potentials

$$V(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0} \frac{qc}{(z c - \mathbf{z} \cdot \mathbf{v})}$$

and

$$\mathbf{A}(\mathbf{r}, t) = \frac{\mu_0}{4\pi} \frac{qc\mathbf{v}}{(z c - \mathbf{z} \cdot \mathbf{v})} = \frac{\mathbf{v}}{c^2} V(\mathbf{r}, t)$$

for a point charge moving with constant velocity v are no longer dependent on the retarded variables \mathbf{z} and t_{ret} . Here $\mathbf{z} = \mathbf{r} - \mathbf{w}(t_{\text{ret}})$ with $\mathbf{w}(t)$ being the position of the point charge q at time t and (see Figure 1).

Exercise 3

In Bohr's theory of hydrogen, the electron in its ground state was supposed to travel in a circle of radius 5 \AA , held in orbit by the Coulomb attraction of the proton. According to classical electrodynamics, this electron should radiate, and hence spiral into the nucleus.

- Show that $v \ll c$ for most of the trip.
- Calculate the lifetime of Bohr's atom assuming that each revolution is essentially circular.